

General information

Phenthoate 50% EC is a systemic organophosphate insecticide with contact and stomach action, is particularly suitable for the control leaf roller; stem borer; borer, scale insects, yanonensis and other pests.

Direction for use

Crop	pests	Rate of use	Application
Rice	Leaf roller; Stem borer; rorer	750-900g/ha	Spraying
Orange tree	Scale insects; Yanonensis; Psyllids	0.625-1g/kg	Spraying

Principle uses

Phenthoate 50%EC is used to control Lepidoptera, jassids, aphids and soft scales; mosquito larvae and adults; blowflies and various strains of houseflies; and ked. It is used to protect cotton, rice, fruit and vegetable crops, and ornamental plants. It is used on animals to protect against external parasites; on and around agricultural structures to control nuisance and disease vector insects.



Phenthoate 50% EC Insecticide

Ingredient	w/v
Phenthoate.....	50%
Inert Ingredients.....	50%
Batch No:	Net contents:
Manuf. date:	Shelf life: 2 YEARS



KEEP OUT OF REACH OF CHILDREN!
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Antidote: atropine

Compatibility:

Incompatible with: strong alkali and acid, and oxidizers.

Protective precautions:

Ingestion: Avoid eating, drinking, smoking or cosmetic application.

Eyes: To avoid eye contact, wear chemical goggles or a full face shield.

Skin: To avoid skin contact, wear full-length disposable protective clothing, waterproof boots, nitrile gloves, and a head covering.

Inhalation: To avoid breathing vapor or mist, wear an approved supplied air respirator or self-contained breathing apparatus.

Poisoning Symptoms:

Perspiration, dizziness, headache, blurred vision, nausea, vomiting, respiratory difficulties, diarrhoea, muscular cramp, salivation.

First aid:

Ingestion: Seek medical assistance.

Eyes: Irrigate exposed eyes with copious amounts of tepid water for at least 15 minutes.

Skin: Systemic effects can occur from dermal exposure to organophosphates. Remove contaminated clothing, wash skin, hair and nails vigorously with repeated soap washings.

Inhalation: Move patient to fresh air. Monitor for respiratory distress. If cough or difficulty breathing develops, evaluate for respiratory tract irritation, bronchitis, or pneumonitis.

Manufacturer:

Shanghai Tenglong Agrochem Co., Ltd.

Address: Yangpu building 24B, No.2005, Huangxing Road, Yangpu Shanghai